

JOURNAL OF FUNCTIONAL ANALYSIS **66**, 67–72 (1986)

A Remark on Compact Automorphism Groups of C^* -Algebras

SERGIO DOPLICHER*

*Dipartimento di Matematica, Università di Roma "La Sapienza,"
I-00185 Roma, Italy*

AND

JOHN E. ROBERTS

*Universität Osnabrück, Fachbereich Physik,
D-4500 Osnabrück, Federal Republic of Germany*

Communicated by A. Connes

Received July 23, 1984

A C^* -system over a compact group whose fixed-point algebra has trivial centre and a relative commutant coinciding with the centre of the full algebra is obtained canonically by inducing up from a C^* -system over a closed subgroup with the same fixed-point algebra but which now has trivial relative commutant. © 1986 Academic Press, Inc.

Let \mathcal{B} be a C^* -algebra with unit, G a compact group, and α a strongly continuous action of G by automorphisms of \mathcal{B} . We shall denote by \mathcal{A} the α_G -fixed-point subalgebra of \mathcal{B} and by \mathcal{C} the centre of \mathcal{B} ,

$$\begin{aligned}\mathcal{A} &\equiv \mathcal{B}^\alpha, \\ \mathcal{C} &\equiv \mathcal{B}' \cap \mathcal{B}.\end{aligned}$$

In this article we show that if the relative commutant of \mathcal{A} in \mathcal{B} is \mathcal{C} and \mathcal{A} has trivial centre,

$$\mathcal{A}' \cap \mathcal{B} = \mathcal{C}, \tag{1}$$

$$\mathcal{A}' \cap \mathcal{A} = \mathbb{C}I, \tag{2}$$

* Research supported by the Ministero della Pubblica Istruzione and CNR-GNAFA.

then there is a C^* -algebra \mathcal{F} and an action τ of a closed subgroup H of G such that \mathcal{A} is the τ_H -fixed points of \mathcal{F} , the relative commutant of \mathcal{A} in \mathcal{F} reduces to the complex numbers and $\{\mathcal{B}, \alpha\}$ is obtained from $\{\mathcal{F}, \tau\}$ by inducing up from the subgroup H to G . This simple construction is used in an essential way to determine the compact Lie group associated with certain endomorphisms of C^* -algebras [1].

If H is a closed subgroup of the compact group G and τ is a strongly continuous action of H by automorphisms of a C^* -algebra \mathcal{F} , the C^* -system $\{\mathcal{C}_\tau(G, \mathcal{F}), \hat{\tau}\}$ obtained from $\{\mathcal{F}, \tau\}$ by inducing up to G is defined as follows. $\mathcal{C}_\tau(G, \mathcal{F})$ is the C^* -algebra of continuous functions X from G to \mathcal{F} (with pointwise $*$ -algebraic operations and supremum norm) satisfying

$$X(hg) = \tau_h(X(g)), \quad h \in H, \quad g \in G; \quad (3)$$

$$\hat{\tau} \text{ is the action of } G \text{ on } \mathcal{C}_\tau(G, \mathcal{F}) \text{ defined by } (\hat{\tau}_g(X))(g_0) = X(g_0 g); \quad g, g_0 \in G, \quad X \in \mathcal{C}_\tau(G, \mathcal{F}). \quad (4)$$

1. THEOREM. *Under assumptions (1) and (2) on the C^* -system $\{\mathcal{B}, \alpha\}$ over the compact group G , there exists a C^* -algebra \mathcal{F} , a closed subgroup H of G , a strongly continuous action τ of H on \mathcal{F} and a monomorphism π of \mathcal{A} into \mathcal{F} such that*

- (a) $\pi(\mathcal{A}) = \mathcal{F}^\tau$,
- (b) $\pi(\mathcal{A})' \cap \mathcal{F} = \mathbb{C}I$,
- (c) $\{\mathcal{B}, \alpha\}$ is isomorphic to $\{\mathcal{C}_\tau(G, \mathcal{F}), \hat{\tau}\}$.

Moreover, H is unique up to conjugation in G and $\{\mathcal{F}, \tau\}$ is then unique up to isomorphism.

The proof will require some elementary lemmas. First, note that (1) and (2) imply $\mathcal{C}^\alpha = \mathbb{C}I$ so that $g \in G \rightarrow \alpha_g|_{\mathcal{C}}$ is an ergodic action of a compact group on a commutative C^* -algebra. The transposed action $g \rightarrow \phi \circ \alpha_g^{-1}$ is then an ergodic action by homeomorphisms of the compact Hausdorff space $\sigma(\mathcal{C})$, the spectrum of \mathcal{C} . As the action is continuous and G is compact, each point in $\sigma(\mathcal{C})$ has a closed orbit so that ergodicity implies that G acts transitively on $\sigma(\mathcal{C})$. If

$$H = \{g \in G \mid \phi \circ \alpha_g = \phi\}$$

is the stabilizer of some chosen $\phi \in \sigma(\mathcal{C})$, H is a closed subgroup of G and $\sigma(\mathcal{C})$ is homeomorphic to the coset space $H \backslash G$.

Let J_ϕ denote the smallest closed two sided (or equivalently left or right) ideal containing $\ker \phi$, J_ϕ is the closed linear span of

$$\mathcal{B} \ker \phi \mathcal{B} = \mathcal{B} \ker \phi = \ker \phi \mathcal{B}.$$

2. LEMMA. *The ideal J_ϕ is the intersection of the kernels of the G.N.S. representations π_ω , where ω ranges over the pure state extensions of ϕ to \mathcal{B} . Moreover*

$$\bigcap_{\phi \in \sigma(\mathcal{C})} J_\phi = \{0\}. \quad (5)$$

Proof. Let \hat{J}_ϕ denote the intersection described in the statement of the lemma. A pure state ω of \mathcal{B} restricts to a pure state of \mathcal{C} , $\omega|_{\mathcal{C}} \in \sigma(\mathcal{C})$, and $\pi_\omega(C) = \omega(C)I$, $C \in \mathcal{C}$. Hence $\ker \pi_\omega \supset \ker(\omega|_{\mathcal{C}})$ so $J_\phi \subset \hat{J}_\phi$. Conversely, every irreducible representation $\tilde{\pi}$ of \mathcal{B}/J_ϕ is of the form $\tilde{\pi}(B + J_\phi) = \pi_\omega(B)$, where ω is a pure state of \mathcal{B} and $\omega|_{\mathcal{C}} = \phi$. Since \mathcal{B}/J_ϕ is separated by its irreducible representations, $\hat{J}_\phi \subset J_\phi$ and equality is proved. We now have

$$\bigcap_{\phi \in \sigma(\mathcal{C})} J_\phi = \bigcap \{ \ker \pi_\omega \mid \omega \text{ is a pure state of } \mathcal{B} \} = \{0\}.$$

If α is an automorphism of \mathcal{B} and $\phi \in \sigma(\mathcal{C})$, then $\alpha(\ker \phi) = \ker \phi \circ \alpha^{-1}$ so that

$$\alpha(J_\phi) = J_{\phi \circ \alpha^{-1}}.$$

In particular, $\alpha_g(J_\phi) = J_{\phi \circ \alpha_g^{-1}}$ so that the stabilizer H of ϕ leaves J_ϕ globally invariant.

Fixing $\phi \in \sigma(\mathcal{C})$, we write

$$J_g = \alpha_g(J_\phi) = J_{\phi \circ \alpha_g^{-1}}, \quad g \in G, \quad (6)$$

for brevity so that, in particular, $J_e = J_\phi$, where e is the unit of G . Let $\eta: \mathcal{B} \rightarrow \mathcal{B}/J_e$ denote the quotient map, $\eta(B) = B + J_e$, $B \in \mathcal{B}$ and let

$$\eta_g = \eta \circ \alpha_g, \quad g \in G, \quad (7)$$

then Eqs. (5) and (6) give

$$\begin{aligned} \ker \eta_g &= \alpha_g^{-1}(J_e) = J_{g^{-1}}, \\ \bigcap_{g \in G} \ker \eta_g &= \{0\}. \end{aligned} \quad (8)$$

The C^* -system $\{\mathcal{F}, \tau\}$ is now defined by

$$\begin{aligned} \mathcal{F} &= \eta(\mathcal{B}), \\ \tau_h \circ \eta &= \eta \circ \alpha_h, \quad h \in H, \end{aligned} \quad (9)$$

and the induced C^* -system $\{\mathcal{C}_\tau(G, \mathcal{F}), \hat{\tau}\}$, introduced above, will be denoted simply by $\{\hat{\mathcal{F}}, \hat{\tau}\}$. Define a homomorphism

$$B \in \mathcal{B} \rightarrow \hat{B} \in \hat{\mathcal{F}} \quad (10)$$

by

$$\hat{B}(g) = \eta_g(B); \quad B \in \mathcal{B}, \quad g \in G.$$

The map (10) is well defined since, for $h \in H, g \in G$, we have by (9),

$$\hat{B}(hg) = \eta_{hg}(B) = \eta \circ \alpha_{hg}(B) = \tau_h \circ \eta \circ \alpha_g(B) = \tau_h(\hat{B}(g)).$$

The map (10) intertwines α and $\hat{\tau}$ since, for $g, g_0 \in G$

$$(\hat{\tau}_g(\hat{B}))(g_0) = \hat{B}(g_0 g) = \eta_{g_0 g}(B) = \eta \circ \alpha_{g_0 g}(B) = \widehat{\alpha_g(B)}(g_0).$$

3. LEMMA. *The map $B \in \mathcal{B} \rightarrow \hat{B} \in \hat{\mathcal{F}}$ defines an isomorphism of $\{\mathcal{B}, \alpha\}$ onto $\{\hat{\mathcal{F}}, \hat{\tau}\}$.*

Proof. By (8) and (10), $\hat{B} = 0$ implies $B = 0$ so we have only to establish that (10) has dense image. Given $X \in \hat{\mathcal{F}}, \varepsilon > 0$, and $g \in G$ there is a $B \in \mathcal{B}$ with

$$\|X(g') - \hat{B}(g')\| < \varepsilon, \quad g' \in \Omega,$$

where Ω is an H -stable neighbourhood of g , i.e., $h\Omega = \Omega, h \in H$. Since G is compact we can find a finite covering $\Omega_1, \Omega_2, \dots, \Omega_n$ of G by H -stable open sets and $B_1, B_2, \dots, B_n \in \mathcal{B}$ with

$$\|X(g) - \hat{B}_i(g)\| < \varepsilon, \quad g \in \Omega_i, \quad i = 1, 2, \dots, n.$$

Pick $C_1, C_2, \dots, C_n \in \mathcal{C}$ so that $\hat{C}_1, \hat{C}_2, \dots, \hat{C}_n$ is a partition of unity by continuous functions constant on left H -cosets subordinate to the covering and let $B = \sum_i C_i B_i$. Then

$$\begin{aligned} \|X(g) - \hat{B}(g)\| &= \left\| \sum_i \hat{C}_i(g)(X(g) - \hat{B}_i(g)) \right\| \\ &\leq \sum_i \hat{C}_i(g) \|X(g) - \hat{B}_i(g)\| < \varepsilon, \quad g \in G, \end{aligned}$$

completing the proof.

Our original argument at this point identified $\hat{\mathcal{F}}$ with the C^* -algebra defined by a continuous field of C^* -algebras over $H \backslash G$ and appealed to the Stone-Weierstrass theorem for continuous fields [2, Corollary 11.5.3]. G.

K. Pedersen kindly drew our attention to a related result [3, Theorem 2.6] which led us to simplify the proof of this lemma.

Proof of the Theorem. Let $\pi: \mathcal{A} \rightarrow \mathcal{F}$ be the restriction of η to the fixed-point algebra \mathcal{A} . By (7)

$$\eta_g(A) = \pi(A), \quad g \in G, \quad A \in \mathcal{A}, \quad (11)$$

so that π is injective by (8). The C^* -systems $\{\mathcal{B}, \alpha\}$ and $\{\hat{\mathcal{F}}, \hat{\tau}\}$ are isomorphic by Lemma 3 so

$$\hat{\mathcal{A}} = \hat{\mathcal{F}}^{\hat{\tau}}. \quad (12)$$

Comparing (10) and (11), $\hat{\mathcal{A}}$ is the set of constant $\pi(\mathcal{A})$ -valued functions on G . On the other hand, $X \in \hat{\mathcal{F}}^{\hat{\tau}}$ if and only if $X(g_0 g) = X(g_0)$, $g, g_0 \in G$, i.e., $X(g) = X(e)$, $g \in G$ and $\tau_h(X(e)) = X(h) = X(e)$, $h \in H$. In other words, $\hat{\mathcal{F}}^{\hat{\tau}}$ is the set of constant \mathcal{F}^{τ} -valued functions on G . So (12) can be read as

$$\pi(\mathcal{A}) = \mathcal{F}^{\tau}.$$

Lemma 3 also gives us the equality

$$(\mathcal{A}' \cap \mathcal{B})^{\wedge} = (\hat{\mathcal{F}}^{\hat{\tau}})' \cap \hat{\mathcal{F}}. \quad (13)$$

Now the restriction of the map $\hat{\cdot}$ to $\mathcal{A}' \cap \mathcal{B}$ is essentially the Gelfand transform: the l.h.s. of (13) is the set of all continuous $\mathbb{C}I$ -valued functions on G constant on left H -cosets. The r.h.s. is the set of all $\pi(\mathcal{A})' \cap \mathcal{F}$ -valued functions in $\hat{\mathcal{F}}$. As the set of values taken by such functions at $e \in G$ is $\pi(\mathcal{A})' \cap \mathcal{F}$, we get

$$\pi(\mathcal{A})' \cap \mathcal{F} = \mathbb{C}I.$$

Thus we have proved (a) and (b) of the Theorem. Part (c) was already proved as Lemma 3 so we are left with the assertions on uniqueness. Let $\{\mathcal{F}_1, H_1, \tau_1, \pi_1\}$ be another choice satisfying (a), (b), and (c) and let $\rho: \mathcal{B} \rightarrow \mathcal{C}_{\tau_1}(G, \mathcal{F}_1)$ be an isomorphism realizing (c). The image of the centre \mathcal{C} of \mathcal{B} under ρ is the centre of $\mathcal{C}_{\tau_1}(G, \mathcal{F}_1)$, i.e., the set of continuous $\mathbb{C}I$ -valued functions on G constant on left H_1 -cosets. Hence there is a $g \in G$ such that

$$\phi(A)I = \rho(A)(g), \quad A \in \mathcal{C}, \quad (14)$$

where $\phi \in \sigma(\mathcal{C})$ is the reference point used to define η . Consequently,

$$H_1 = gHg^{-1}.$$

Modulo an automorphism in α_G , we may now assume that $g = e$ in (14). The homomorphism

$$X \in \mathcal{C}_{\tau_1}(G, \mathcal{F}_1) \rightarrow X(e) \in \mathcal{F}_1$$

with kernel J_1 , say, gives us an exact sequence

$$0 \rightarrow J_1 \rightarrow \mathcal{C}_{\tau_1}(G, \mathcal{F}_1) \rightarrow \mathcal{F}_1 \rightarrow 0.$$

By definition, we have an exact sequence

$$0 \rightarrow J_\phi \rightarrow \mathcal{B} \rightarrow \mathcal{F} \rightarrow 0$$

and, since $g = e$ in (14), $\rho(J_\phi) = J_1$. Hence the diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & J_1 & \longrightarrow & \mathcal{C}_{\tau_1}(G, \mathcal{F}_1) & \longrightarrow & \mathcal{F}_1 & \longrightarrow & 0 \\ & & \uparrow \rho & & \uparrow \rho & & \uparrow \rho_1 & & \\ 0 & \longrightarrow & J_\phi & \longrightarrow & \mathcal{B} & \longrightarrow & \mathcal{F} & \longrightarrow & 0 \end{array}$$

defines an isomorphism ρ_1 of \mathcal{F} onto \mathcal{F}_1 . Since ρ intertwines the actions α and $\hat{\tau}_1$ of G and J_ϕ and J_1 are stable under the restriction of these actions to H , ρ_1 must intertwine the actions τ and τ_1 of H .

ACKNOWLEDGMENTS

S. D. would like to thank Professors D. Buchholz, R. Haag, and L. Streit for the warm hospitality at the University of Hamburg and at ZiF, University of Bielefeld. J. E. R. gratefully acknowledges the support extended by the Department of Mathematics of the University of Rome "La Sapienza."

REFERENCES

1. S. DOPLICHER AND J. E. ROBERTS, Compact Lie groups associated with endomorphisms of C^* -algebras, *Bull. Amer. Math. Soc.* **11** (1984), 333–348.
2. J. DIXMIER, " C^* -Algebras," North-Holland, Amsterdam/New York/Oxford, 1977.
3. D. OLESEN AND G. K. PEDERSEN, Applications of the Connes spectrum to C^* -dynamical systems, III, *J. Funct. Anal.* **45** (1982), 357–390.